

FORAGE AND LIVESTOCK RESEARCH - 1988

RESEARCH CENTER TECHNICAL REPORT 88-1

Texas A&M University Agricultural Research & Extension Center
at Overton

Texas Agricultural Experiment Station
Texas Agricultural Extension Service

Overton, Texas

April 21, 1988

All programs and information of the Texas Agricultural Experiment Station and Texas Agricultural Extension Service are available to everyone without regard to race, color, religion, sex, age, or national origin.

Mention of trademark or a proprietary product does not constitute a guarantee or a warranty of the product by the Texas Agricultural Experiment Station or Texas Agricultural Extension Service and does not imply its approval to the exclusion of other products that also may be suitable.

COASTAL BERMUDAGRASS RESPONSE TO FLUID COMBINATIONS OF NITROGEN,
PHOSPHORUS, POTASSIUM, AND MAGNESIUM AND METHODS OF APPLICATION

V. A. Haby, J. V. Davis, and A. T. Leonard

SUMMARY

Blends of nitrogen (N), phosphorus (P), potassium (K), and magnesium (Mg) were made with urea-ammonium nitrate (32-0-0), ammonium polyphosphate (11-37-0), muriate of potash (0-0-60), and magnesium chloride (0-0-0-8.1% Mg). Pounds of nutrients applied per acre for each flush of Coastal bermudagrass growth as individual treatments were 80-0-0, 80-22-0, 80-0-60, 80-22-60, and 80-22-60-12. Each treatment was applied as a broadcast spray, surface dribble banded, and subsurface dribble banded. Coastal bermudagrass dry matter production generally increased due to P fertilization the second year on the Lilbert soil site. Potassium began to provide significant yield increases the third year on the Gallime soil site. Surface dribble banding these solutions increased yields the second year on the Lilbert soil and the first year on the Gallime soil. In most years, method of application had no significant effect on forage production on these acid, sandy soils.

INTRODUCTION

Urea-ammonium nitrate solutions, as with solid fertilizers, can be combined into blends with phosphorus (P), potassium (K), magnesium (Mg), and other soluble fertilizer materials. East Texas soils are generally deficient in P and K, and Mg has increased grass yield in at least one research trial at Overton. This research was designed to evaluate the combinations of UAN with P, K, and Mg applied as broadcast spray, surface dribble bands, and subsurface dribble bands. The main objective of this study was to determine Coastal bermudagrass response to these fertilizer nutrients on two soils, and to evaluate forage response to methods of application of the blends of N, P, K, and Mg.

PROCEDURE

Fluid blends of N, P, K, and Mg were made with urea-ammonium nitrate (UAN, 32-0-0), ammonium polyphosphate (11-37-0), muriate of

potash (0-0-60), and magnesium chloride (0-0-0-8.1% Mg). These blends were applied to acid, sandy Lilbert (initial pH 5.0) and Gallime (pH 5.0) soils. Pounds of nutrients applied as N-P₂O₅-K₂O-Mg/ac for each flush of grass growth were 80-0-0, 80-22-0, 80-0-60, 80-22-60, and 80-22-60-12. Each treatment was applied spray broadcast, or surface dribble banded, or subsurface dribble banded using a coulter and back-swept knife. Spacings between dribble bands were 14 inches. These solutions were applied to 10 x 20 foot plots. Harvests were made after first trimming one foot off the end of each plot with a Hege 211B forage plot harvester. A 4.9 foot center swath was harvested from the length of each plot and was weighed. A dry matter sample was collected from each plot for moisture and chemical analysis. Yield data were analyzed statistically by MSUSTAT for analysis of variance and Newman-Keuls mean comparisons.

RESULTS

Yields were not increased at either site by fertilization with P, K, or Mg the first year of the study (Table 1). The blend of N and P

TABLE 1. RESPONSE OF COASTAL BERMUDAGRASS TO COMBINATIONS OF PHOSPHORUS, POTASSIUM, AND MAGNESIUM WITH UAN, AND TO METHODS OF APPLICATION OF THESE SOLUTIONS TO LILBERT AND GALLIME SOILS

Nutrient Blend ²	Dry Matter Yield ¹					
	Lilbert Soil			Gallime Soil		
	1984	1985	1986	1984	1985	1986
N-P ₂ O ₅ -K ₂ O-Mg						
----lb/ac----	-----Tons/ac-----					
80- 0- 0- 0	3.9 a	3.8 a	4.9 a	6.3 c	6.1 a	4.8 a
80-22- 0- 0	3.9 a	5.0 c	6.2 bc	6.1 bc	6.6 ab	5.4 a
80- 0- 60- 0	3.4 a	4.2 ab	5.5 ab	5.3 a	6.5 ab	6.8 b
80-22- 60- 0	3.8 a	4.9 bc	7.0 d	5.7 ab	6.9 ab	7.7 b
80-22- 60- 12	3.9 a	4.9 bc	6.7 cd	6.1 bc	7.6 b	7.5 b
<u>Applic. Method</u>						
Spray Broadcast	3.7 a	4.5 ab	5.9 a	5.5 a	6.6 a	6.5 a
Surf. Drib. Bd.	3.8 a	4.9 b	6.3 a	6.0 b	7.0 a	6.6 a
Subsurf. " "	3.8 a	4.3 a	6.0 a	6.2 b	6.8 a	6.2 a

¹Dry matter yields within an individual year and site, by solution blends or by method of application, followed by a similar letter are not significantly different at p<.05 level of probability by Newman-Keuls mean comparisons.

²Actual rates of N, P₂O₅, K₂O, and Mg at each application.

significantly increased dry matter yield on the Lilbert soil the second year. A trend indicates that P and K blended with N increased yields on the Gallime soil, but a significant yield increase occurred when Mg was added with N, P, and K compared to N alone. The combination of N, P, and K significantly increased dry matter production on the Lilbert soil the third year. Any nutrient combination that included K significantly increased forage production the third year on the Gallime soil. Both research sites were previously in poorly fertilized hay meadow-pasture situations prior to initiation of this study. There was apparently sufficient nutrient recycling occurring to provide a small reserve of soil P, K, and Mg for one or two years of intensive forage production.

Surface dribble band application of these nutrient solutions improved dry matter yield compared to subsurface dribble banding on the Lilbert soil the second year, and slightly increased yield compared to the spray broadcast application. Dribble banding on the Gallime soil significantly improved dry matter production compared to spray broadcast application the first year. It is noteworthy that banding fluid blends of N, P, K, and Mg improved dry matter yield; whereas, banding UAN solution alone did not improve yields compared to the spray broadcast application. It may be possible that concentrating the P in a band limited its contact with the soil, thereby preventing its precipitation as insoluble aluminum phosphates, allowing the P to remain available to the plant.